

## REPTILIA: TESTUDINES: CHELIDAE


## PLATEMYS, PLATEMYS PLATYCEPHALA

## Catalogue of American Amphibians and Reptiles.

ERNST, CARL H. 1987. *Platemys, Platemys platycephala*.***Platemys* Wagler  
Twistnecked turtles***Platemys* Wagler, 1830:135. Type-species, *Testudo planiceps* Schoepff, 1801:115 (= *Testudo platycephala* Schneider, 1792:261) by monotypy.*Emys*: Gray, 1831:40. In part.*Hydraspis*: Gray, 1831:40. In part.

- CONTENT. One species, *Platemys platycephala*.

● DEFINITION. Adult males grow to 18 cm carapace length, adult females to 16.5 cm. The flat, elliptical carapace has a deep medial groove flanked by two longitudinal keels. The carapace is highest just anterior to the 2nd intervertebral seam and widest at level of the 7th or 8th marginals. The posterior marginals are flared, those lateral are upturned. Neural bones are usually absent; the 8 pairs of costal bones meet at the midline. The nuchal extends to the carapacial rim; there are 11 pairs of peripherals. The yellow carapace has varying amounts of dark pigment. The hingeless plastron is dark brown with a yellow border; the yellow bridge has a dark transverse bar. The forelobe is slightly upturned, and is longer and slightly wider than the hindlobe which is posteriorly notched. The intergular separates the gulars and is about half as long as the plastral forelobe. Both axillary and inguinal buttresses are strong; the inguinals are ankylosed to the 5th costal bones. The broad, hexagonal-shaped entoplastron lies anterior to the humeropectoral seam. The flat, broad skull is extensively roofed and parallel-sided. The prefrontals do not meet at the midline. The moderate temporal arch is formed by the squamosal and parietals. The parietal does not touch the supraoccipital. The vomer contacts the parietals and palatines; the palatines are not reduced. The unnotched jaws are ridgeless. Dorsally the head is covered with smooth, undivided skin, laterally there are 1-3 rows of large scales. The chin has two barbels. The head is orange to yellowish dorsally, dark brown laterally and ventrally; the light dorsal pigment extends downward to midway between the orbit and tympanum. The 5th and 8th cervical centra are biconvex (Williams, 1950). The neck has numerous blunt



MAP. Solid circles mark the type-localities; open circles other selected localities.

tubercles dorsolaterally; the thighs bear small blunt tubercles. The toes are webbed and the anterior surfaces of the limbs have large scales. Males have concave plastra and longer, thicker tails; females have the posterior tips of the plastron bent upward.

- DESCRIPTIONS, ILLUSTRATIONS, AND DISTRIBUTION. See species account.

● FOSSIL RECORD. Several late Cretaceous to late Pleistocene southern South American turtle fossils have been assigned to *Platemys*: *P. antiqua*, *P. fossilis*, *P. laevius*, *P. robusta* (Ameghino, 1880); *P. paranensis* (Bravard, 1883); *P. torrentium* (Burmeister, 1885); *P. bergii*, *P. enteriana*, *P. holmbergii*, *P. robusta*, *P. mesopotamica* (Ambrosetti, 1893); *P. quarantinica*, *P. patagonica*, and *P. sehuensis* (Ameghino, 1899). However, Wood (in prep.) has found all of these to be either *nomina nuda* or *nomina dubia*; so, at present, no fossils can be accurately attributed to this genus.

- PERTINENT LITERATURE. See species account.

● ETYMOLOGY. The generic name *Platemys* is from the Greek *platos*, flat, and *emydos*, a freshwater turtle, and refers to the depressed carapace.

***Platemys platycephala* (Schneider)  
Twistnecked turtle**

*Testudo platycephala* Schneider, 1792:261. Type-locality, "ost-Indien," restricted to "Cayenne, French Guiana" by Ernst, 1984:350. Holotype, undesignated.

*Testudo planiceps* Schneider, 1792:pl. 7. Type-locality, not stated, restricted to "Cayenne," French Guiana by Martin (in Schweigger, 1812). Holotype, undesignated.

*Testudo planiceps* s. *platycephala*: Schoepff, 1801:115.

*Testudo martinella* Daudin, 1803:344. Type-locality, "Cayenne," French Guiana from label. Holotype, Mus. Nat. Hist. Natur., Paris 8760, mounted adult female, collector and date unknown (examined by author).

*Emys discolor* Thunberg, in Schweigger, 1812:302. Type-locality, unknown, restricted to "Cayenne, French Guiana" by Ernst, 1984:350. Holotype, undesignated.

*Emys planiceps*: Schweigger, 1812:303.

*Testudo dicolor*: Ekstrand, 1823:6. Ex errore.

*Emys canaliculata* Spix, 1824:10. Type-locality, "subcognomine Japutipirema in campus paludosis juxta ripam fluminis Solimoens," Brazil. Holotype, Zool. Staatssammlung München 3007/0, shell and skeleton, probably collected by Spix or Martius in 1819 (examined by author).

*Hydraspis planiceps*: Bell, 1828:512.

*Hydraspis canaliculata*: Bell, 1828:512.

*Clemmys planiceps*: Ritgen, 1828:272.

*Emys platicephala*: Cuvier, 1829:11. Ex errore.

*Emys carunculata*: Cuvier, 1829:11. Ex errore.

*Platemys planiceps*: Wagler, 1830:135.

*Emys caniculata*: Gray, 1831:40. Ex errore.

*Hydraspis constricta* Gray, 1831:43. Type-locality, unstated, re-



FIGURE 1. *Platemys platycephala*, dorsolateral view. Photograph by Roger W. Barbour.



FIGURE 2. *Platemys platycephala*, ventrolateral view. Photograph by Roger W. Barbour.

stricted to "Belem, Brazil" by Ernst, 1984:350. Holotype, undesigned.

*Emys martinella*: Schinz, 1833:45. Ex errore.

*Platemys Martinella*: Duméril and Bibron, 1835:407.

*Hydraspis discolor*: Fitzinger, 1835:126.

*Platemys platycephala*: Boulenger, 1889:227. First use of combination.

*Platemys platycephala*: Kasper, 1903:161. Ex errore.

*Platemys platemys*: Dunn, 1945:327. Ex errore.

*Platemys platycephala*: Orcés, 1949:21. Ex errore.

● CONTENT. Two subspecies: *Platemys platycephala platycephala* and *P. p. melanonota*.

● DEFINITION. See generic account.

● DESCRIPTIONS. General descriptions: Spix (1824), Gray (1855), Boulenger (1889), Siebenrock (1909), Luederwaldt (1926), Freyete (1977), Duellman (1978), Pritchard (1967, 1979), Ernst (1984), and Pritchard and Trebbau (1984). Other descriptions: vertebrae (Kasper, 1903; Williams, 1950); coracoid (Walker, 1973); pubis (Grassé, 1970); skull (Gaffney, 1977, 1979); digestive tract (Jacobshagen, 1920; Parsons and Cameron, 1976); rostral pores (Winokur and Legler, 1974); ear (Wever, 1978); karyotype (Gorman, 1973; Barros et al., 1976; Bull and Legler, 1980; McBee et al., 1985).

● ILLUSTRATIONS. Color illustrations of adults occur in Pritchard (1967, 1979) and Pritchard and Trebbau (1984); other illustrations of adults are in Mertens (1954), Wermuth and Mertens (1961), Freyete (1977), Freiberg (1981) and Ernst (1984). The skull was illustrated by Wermuth and Mertens (1961), and Gaffney (1977, 1979); the vertebrae by Kasper (1903); the karyotype by Barros et al. (1976), Bull and Legler (1980), and McBee et al. (1985), and the egg and juvenile by Medem (1983).

● DISTRIBUTION. *P. platycephala* occurs in the Caribbean drainages of eastern Venezuela and the Guianas, and in the Amazon Basin from northeastern Bolivia, eastern Ecuador and Perú, south-eastern Colombia and southern Venezuela eastward to Belem, Brazil.

● FOSSIL RECORD. See generic account.

● PERTINENT LITERATURE. General accounts are in Goeldi (1905), Mertens (1954), Freyete (1977), Pritchard (1964, 1967, 1979), Freiberg (1972, 1981), and Pritchard and Trebbau (1984). Additional references are listed by topic. Reproduction and embryology: Medem (1960, 1983), Carillo de Espinoza (1970), Dixon and Soini (1977), Ewert (1979, 1985), Harding (1983), Pickering (1984). Anatomy: Stannius (1856), Parsons (1968). Hearing: Wever (1978). Blood: Sullivan and Riggs (1967). Longevity: Bowler (1977). Zoogeography: Ernst (1984). Vernacular names: Mittermeier et al. (1980). Morphometry: Ernst and Lovich (1986). Distribution: Luederwaldt (1926), Medem (1957), Hoogmoed (1979). Taxonomic relationships: Ernst (1984), Pritchard (1984), McBee et al. (1985). Disease: Jacobson et al. (1982).

● ETYMOLOGY. The specific name *platycephala* is from the Greek *platos*, flat, and *kephale*, head, referring to the flat crown. The subspecific name *melanonota* is from the Greek *melania*, blackness, and *notos*, back, referring to the dark carapace.

## 1. *Platemys platycephala platycephala* (Schneider)

*Testudo platycephala* Schneider, 1792:261. See species synonymy.

*Testudo planiceps* Schneider, 1792:pl. 7. See species synonymy.

*Testudo planiceps* s. *platycephala*: Schoepff, 1801:115.

*Testudo martinella* Daudin, 1803:344. See species synonymy.

*Emys discolor* Thunberg, in Schweigger, 1812:302. See species synonymy.

*Emys planiceps*: Schweigger, 1812:303.

*Testudo discolor*: Ekstrand, 1823:6.

*Emys canaliculata* Spix, 1824:10. See species synonymy.

*Hydraspis planiceps*: Bell, 1828:512.

*Hydraspis canaliculata*: Bell, 1828:512.

*Clemmys planiceps*: Ritgen, 1828:272.

*Emys platycephala*: Cuvier, 1829:11.

*Emys carunculata*: Cuvier, 1829:11.

*Platemys planiceps*: Wagler, 1830:135.

*Emys caniculata*: Gray, 1831:40.

*Hydraspis constricta* Gray, 1831:43. See species synonymy.

*Emys martirella*: Schinz, 1833:45.

*Platemys Martinella*: Duméril and Bibron, 1835:407.

*Hydraspis discolor*: Fitzinger, 1835:126.

*Platemys platycephala*: Boulenger, 1889:227.

*Platemys platycephala*: Kasper, 1903:161.

*Platemys platemys*: Dunn, 1945:327.

*Platemys platycephala*: Orcés, 1949:21.

*Platemys platycephala platycephala*: Ernst, 1984:347. First use of combination.

● DEFINITION. The dark carapacial pigment is restricted to the borders of the seams between the vertebrae and pleurals and to an incomplete band extending downward through the 2nd and 3rd pleurals to the carapacial rim; the dark bar crosses less than 80% of the bridge; the intergular is wide ( $\bar{x}$  = 27% of anterior plastral width); and the seam separating the intergular and humeral scutes is long ( $\bar{x}$  = 21.7% of anterior plastral width). Usually there are more than 4 postorbital scales and less than 4 scales between the orbit and the tympanum.

## 2. *Platemys platycephala melanonota* Ernst

*Platemys platycephala melanonota* Ernst, 1984:347. Type-locality, "vicinity of Galilea, on the Río Santiago, Amazonas, Perú (4°1'S, 77°47'W)." Holotype, U.S. Nat. Mus. 224136, adult male, collected 1 February 1980 by Roy W. McDiarmid.

● DEFINITION. A dark subspecies with the light carapacial pigment restricted to the vertebral groove, anterior of 1st pleurals, and posterior of 4th pleurals; with a dark bar crossing more than 90% of bridge; with a narrow intergular ( $\bar{x}$  = 24% of anterior plastral width); and with the seam separating the intergular and humeral scutes being relatively short ( $\bar{x}$  = 18.8% of anterior plastral width). Usually there are less than 4 post-orbital scales and more than 3 scales between the orbit and tympanum.

### COMMENT

Schneider (1792) used two names, *Testudo platycephala* and *T. planiceps*, for this species in the original publication; and Schoepff (1801), as first reviewer, apparently gave preference to *planiceps*. *Planiceps* was in general use until Boulenger (1889) selected *platycephala* for the specific name. Thus, for almost 100 years, *platycephala* has been accepted, and due to this customary usage should be retained as the official name (see Pritchard and Trebbau, 1984, for a detailed discussion).

The description of *Hydraspis constricta* by Gray (1831:43) is vague, and since no type-specimen was designated, the allocation of this name as a synonym of *Platemys platycephala* is arbitrary.

The living species *macrocephala*, *pallidipectoris*, *radiolata*, and *spixii* are usually assigned to *Platemys*, but McBee et al. (1985) have discovered significant differences between their karyotypes and that of *platycephala*; Rhodin and Mittermeier (in prep.) have found several skeletal differences between these species and *platycephala* and are preparing a manuscript that will allocate them to a new genus.

### LITERATURE CITED

Ambrosetti, Juan B. 1893. Contribucion al estudio de las tortugas fluviales Oligocenas de los terrenos Terciarios antiguos del Paraná. Bol. Inst. Geogr. Argentina 14:489-499.

- Ameghino, Florentino. 1880. La formación pampeana. Paris. 370 p.
- 1899. Sinopsis geológico-paleontológica. Suplemento (Adiciones y correcciones). La Plata, 13 p.
- Barros, R. M., M. M. Sampaio, M. F. Assis, M. Ayres, and O. R. Cunha. 1976. General considerations on the karyotypic evolution of *Chelonia* from the Amazon region of Brazil. *Cytologia* 41:559-565.
- Bell, Thomas. 1828. Characters of the order, families, and genera of the Testudinata. *Zool. J.* 3:513-516.
- Boulenger, George A. 1889. Catalogue of the chelonians, rhynchocephalians, and crocodiles in the British Museum (Natural History). Taylor and Francis, London. x + 311 p.
- Bowler, J. Kevin. 1977. Longevity of reptiles and amphibians in North American collections as of 1 November, 1975. *Soc. Stud. Amphib. Rept., Herpetol. Circ.* (6), iv + 32 p.
- Bravard, Augusto. 1883-1891. Monografía de los terrenos marinos Terciarios, de las cercanías del Paraná. Buenos Aires Mus. Nac. An. 3:45-94.
- ✓ Bull, James L., and John M. Legler. 1980. Karyotypes of side-necked turtles (Testudines: Pleurodira). *Canadian J. Zool.* 58(5): 828-841.
- Burmeister, Hermann C. C. 1885. Exámen crítico de los mamíferos y reptiles fósiles denominados por D. Augusto Bravard y mencionados en su obra precedente. *An. Mus. Nac. Buenos Aires* 3:95-174.
- ✓ Carillo de Espinoza, Nelly. 1970. Contribución al conocimiento de los reptiles del Perú. *Publ. Mus. Hist. Natur. Javier Prado Ser. A., Zool.* (22):1-64.
- Cuvier, G. L. C. F. D. 1829. *Le regne animal* . . . Nouvelle édition. Paris. Volume 2.
- Daudin, F. M. 1803. Histoire naturelle generale et particulière des reptiles. Vol. 8. F. Dufart, Paris. 439 p.
- Dixon, James R., and Pekka Soini. 1977. The reptiles of the upper Amazon basin, Iquitos region, Peru. II. Crocodylians, turtles and snakes. *Contr. Biol. Geol., Milwaukee Publ. Mus.* (12):1-91.
- Duellman, William E. 1978. The biology of an equatorial herpetofauna in Amazonian Ecuador. *Univ. Kansas Mus. Natur. Hist. Misc. Publ.* (65):1-352.
- Duméril, A. M. C., and G. Bibron. 1835. *Erpétologie générale ou histoire naturelle complète des reptiles*. Vol. 2. Librairie Encyclopédique de Roret, Paris. iv + 682 p.
- Dunn, Emmet R. 1945. Los géneros de anfibios y reptiles de Colombia (IV). *Caldasia* 3(13):308-335.
- Ekstrand, C. H. 1823. *D. D. Fauna brasiliensis quam venia exp. facult. med. Upsala praeside C. P. Thunberg*. Upsala: 10 p.
- ✓ Ernst, Carl H. 1984. Geographic variation in the neotropical turtle *Platemys platycephala*. *J. Herpetol.* (1983) 17(4):345-355.
- ✓ —, and Jeffrey E. Lovich. 1986. Morphometry in the chelid turtle, *Platemys platycephala*. *The Herpetological Journal* 1(2):66-70.
- Ewert, Michael A. 1979. The embryo and its egg: development and natural history, p. 333-413. *In* M. Harless, and H. Morlock (eds.), *Turtles: perspectives and research*. John Wiley & Sons, New York.
- 1985. Embryology of turtles, p. 75-267. *In* C. Gans, F. Billett, and P. F. A. Maderson (eds.), *Biology of the Reptilia*. Vol. 14. John Wiley & Sons, New York.
- Fitzinger, L. J. F. J. 1835. Entwurf einer systematischen Anordnung der Schildkröten nach den Grundsätzen der natürlichen Methode. *Ann. Mus. Wien* 1:103-128.
- Freiberg, Marcos. 1972. Las tortugas (Testudines), p. 583-620. *In* L. Cendrero (ed.), *Zoología Hispanoamericana*. Vertebrados. Editorial Porrúa, S. A., México, D. F.
- ✓ — 1981. Turtles of South America. T.F.H. Publ., Inc., Neptune, New Jersey. 125 p.
- ✓ Freyete, Jacques. 1977. Les chéloniens de Guyane française. I. Etude préliminaire. *Mém. Univ. Paris VI*, 202 p.
- Gaffney, Eugene S. 1977. The side-necked turtle family Chelidae: a theory of relationships using shared derived characters. *Amer. Mus. Novitates* (2620):1-28.
- 1979. Comparative cranial morphology of Recent and fossil turtles. *Bull. Amer. Mus. Natur. Hist.* 164(2):65-376.
- Goeldi, Emilio. 1905. Chelonios do Brasil (Jaboty-Kágados Tartarugas). Capítulo primeiro da monographia "Reptis do Brazil" (Obra inedita, escripta entre 1892-1894). *Bol. Mus. Goeldi, Para, Brazil* 4(4):699-756.
- Gorman, George C. 1973. The chromosomes of the Reptilia, a cytotaxonomic interpretation, p. 350-424. *In* A. B. Chiarelli, and E. Capanna (eds.), *Cytotaxonomy and vertebrate evolution*. Academic Press, London.
- Grasse, Pierre-P. 1970. *Traité de zoologie. Anatomie, systématique biologique*. XIV. Reptiles caracteres générales et anatomie. i-xxxiii + 680 p.
- Gray, John E. 1831. *Synopsis reptilium*. Part 1. *Cataphracta*. Tortoises, crocodiles, and enaliosaurians. Treuttal, Wurtz, and Co., London. viii + 85 p.
- 1855. Catalogue of shield reptiles in the collection of the British Museum. Part 1. Testudinata (tortoises). Taylor and Francis, London. 79 p.
- ✓ Harding, James H. 1983. *Platemys platycephala* (twistneck turtle) reproduction. *Herp. Review* 14(1):22.
- Hoogmoed, Marinus S. 1979. The herpetofauna of the Guanan Region, p. 241-279. *In* W. E. Duellman (ed.), *The South American herpetofauna: its origin, evolution, and dispersal*. *Monogr. Mus. Nat. Hist. Univ. Kansas* (7).
- Jacobson, E. 1920. Zur Morphologie des Oberflächenreliefs den Rumpfdarmschleimhaut des Reptilien. *Z. Naturw. Jena* 56:361-430.
- Jacobson, Elliott R., Jack M. Gaikin, Susan Clubb, and Maron B. Calderwood. 1982. Papilloma-like virus infection in Bolivian side-neck turtles. *J. Am. Veterin. Med. Assoc.* 181(11):1325-1328.
- Kasper, Ad. 1903. Ueber den Atlas und Epistropheus bei den pleurodiren Schildkröten. *Arbeit. Zool. Inst. Wien.* 14:137-172.
- Luederwaldt, Hermann. 1926. Os chelonios Brasileiros. Com a lista das especies do Museu Paulista. *Rev. Mus. Paulista.* 14: 404-468.
- ✓ McBee, Karen, John W. Bickham, Anders G. J. Rhodin, and Russell A. Mittermeier. 1985. Karyotypic variation in the genus *Platemys* (Testudines: Pleurodira). *Copeia* 1985(2):445-449.
- Medem, Federico. 1957. Informe sobre Reptiles Colombianos (II). El conocimiento actual sobre la distribución geográfica de las Testudinata en Colombia. *Bol. Mus. Ciencias Natur.* (2-3): 13-45.
- 1960. Datos zoo-geográficos y ecológicos sobre los Crocodylia y Testudinata de los rios Amazonas, Putumayo y Caquetá. *Caladasia* 8(38):341-351.
- 1983. Reproductive data on *Platemys platycephala* (Testudines: Chelidae) in Colombia, p. 429-434. *In* A. G. J. Rhodin and K. Miyata (eds.), *Advances in herpetology and evolutionary biology. Essays in honor of Ernest E. Williams*. *Mus. Comp. Zool., Cambridge, Massachusetts*.
- Mertens, Robert. 1954. Bemerkenswerte Schildkröten aus Süd- und Zentralamerika. *Aquar.-Terrar. Zeitschr.* 7(9):239-242.
- Mittermeier, Russell A., Federico Medem, and Anders G. J. Rhodin. 1980. Vernacular names of South American turtles. *Soc. Stud. Amphib. Rept., Herpetol. Circ.* (9) 44 p.
- Orcés V., Gustavo. 1949. Los Testudinata ecuatorianos que se conservan en las colecciones de Quito, Ecuador (con excepción de las especies de Galápagos). *Bol. Inform. Cient. Nac., Quito.* 3(20-21):13-22.
- Parsons, Thomas S. 1968. Variation in the choanal structure of Recent turtles. *Canadian J. Zool.* 46(6):1235-1263.
- , and John E. Cameron. 1976. Internal relief of the digestive tract, p. 159-223. *In* C. Gans, and T. S. Parsons (eds.), *Biology of the Reptilia*, vol. 6. Academic Press, London.
- Pickering, Darell. 1984. Twist-neck turtle hatched at Tulsa Zoological Park. *Delaware Herpetol. Soc. Newsletter* 3:1.
- Pritchard, Peter C. H. 1964. Turtles of British Guiana. *Proc. British Guiana Mus. Zool.* (39):19-45.
- ✓ — 1967. *Living turtles of the world*. T.F.H. Publ., Inc., Jersey City, New Jersey. 288 p.
- ✓ — 1979. *Encyclopedia of turtles*. T.F.H. Publ., Inc., Neptune, New Jersey. 895 p.
- 1984. Piscivory in turtles, and evolution of the long necked Chelidae, p. 87-110. *In* M. W. J. Ferguson (ed.), *The structure, development and evolution of reptiles*. *Symp. Zool. Soc. London* (52). Academic Press, London.
- ✓ —, and Pedro Trebbau. 1984. The turtles of Venezuela. *Soc. Stud. Amphib. Rept. Contrib. Herpetol.* (2): viii + 403 p.
- Rhodin, Anders G. J., and Russell A. Mittermeier. *In prep.* Generic revision of the South American chelid turtle genus *Platemys*.
- Rütgen, F. A. 1828. Versuch einer natürlichen Eintheilung der

- Amphibien. *Nova Acta Physico-Medica Acad. Caes. Leopold Carol. Natur. Curio.* 14:246-284.
- Schinz, Heinrich R. 1833-1835. *Naturgeschichte und Abbildungen der Reptilien.* Brodmanns Lithogr., Leipzig. 240 + IV p.
- Schneider, J. G. 1792. *Beschreibung und Abbildung einer neuen Art von Wasserschilddröte.* *Schriften Ges. Naturf. Freunde Berlin* 10:259-284.
- Schoepff, I. D. 1792-1801. *Historia testudinvm uncuribus illustrata.* Ioannis Iacobi Palm., Erlange. xii + 136 p.
- Schweigger, A. F. 1812. *Prodromus monographiae cheloniorum.* *Königsberger Archiv. Naturwiss. Math.* 1:271-368, 406-468.
- Siebenrock, Friedrich. 1909. *Synopsis der rezenten Schildkröten.* *Zool. Jahrb., Suppl.* 10:427-618.
- Spix, Johann Baptist de. 1824. *Animalia nova, sive species novae testudinum et ranarum, quas in itinere per Brasiliam annis MDCCCXVII-MDCCCXX jussu et auspiciis Maximiliana Josephi I. Franc. Seraph. Hubschmanni, Monachii.* 24 p.
- Stannius, Hermann. 1856. *Zweites Buch. Die Amphibien,* p. 1-270. *In P. F. von Siebold and H. Stannius (eds.), Handbuch der Zoologie.* Verlag von Veit & Co., Berlin.
- Sullivan, Bolling, and Austen Riggs. 1967. *Structure, function and evolution of turtle hemoglobins. I. Distribution of heavy hemoglobins. II. Electrophoretic studies.* *Comp. Biochem. Physiol.* 23:437-458.
- Wagler, Johann. 1830. *Natürliches system der amphibien, mit vorangehender Classification der Säugethiere und Vögel.* J. G. Cotta'schen Buchhandlung, München, Stuttgart und Tübingen. vi + 354 p.
- Walker, Warren F. 1973. *The locomotor apparatus of Testudines,* p. 1-100. *In C. Gans, and T. S. Parsons (eds.), Biology of the Reptilia, Vol. 4.* Academic Press, London.
- Wermuth, Heinz, and Robert Mertens. 1961. *Schildkröten, Krokodile, Brückenechsen.* G. Fischer, Jena. 422 p.
- Wever, E. G. 1978. *The reptile ear.* Princeton Univ. Press, Princeton, New Jersey. xii + 1024 p.
- Williams, Ernest E. 1950. *Variation and selection in the cervical central articulations of living turtles.* *Bull. Amer. Mus. Natur. Hist.* 94(9):507-561.
- Winokur, Robert M., and John M. Legler. 1974. *Rostral pores in turtles.* *J. Morphol.* 143(1):107-120.
- Wood, Roger C. *In prep.* *A review of the fossil Chelidae (Testudines, Pleurodira) of South America, with a description of several new occurrences.*

C. H. ERNST, GEORGE MASON UNIVERSITY, FAIRFAX, VIRGINIA 22030.

Primary editor for this account, Jaime D. Villa.

Published 27 March 1987. Copyright 1987 by the SOCIETY FOR THE STUDY OF AMPHIBIANS AND REPTILES.